

SPIN ALIGNMENT OF VECTOR MESON IN HIGH ENERGY REACTIONS

QING-HUA XU, CHUN-XIU LIU*and ZUO-TANG LIANG

Physics Department, Shandong University, Ji'nan, Shandong 250100, China

The recent data on the polarization of vector meson at LEP show that the vector mesons favor the helicity zero state. We calculate the helicity density matrix of vector meson which contains a polarized fragmenting quark by adding the spin of the fragmenting quark and that of the antiquark created in the fragmentation. The data at LEP imply a significant polarization for the antiquark in the opposite direction as that of the fragmenting quark. We extend the calculations to other reactions and make predictions for vector mesons in deeply inelastic lepton-nucleon scatterings and polarized pp collisions.

1. Introduction

Recently, the OPAL and DELPHI collaborations measured the polarizations of vector mesons in e^+e^- annihilation at Z^0 pole at LEP. The data¹ show clearly that the vector mesons in particular those with large momentum fraction favor the helicity zero state. Obviously, the study of this subject should provide us important information on the hadronization mechanism especially the spin effects in fragmentation processes. Compared with hyperons, the study of vector meson polarization has two advantages: First, the decay contribution of heavier hadrons to vector meson production is much smaller. Second, the production rate of vector meson is in general higher than that of hyperon. It is thus interesting and important to ask what we can learn from the data about the polarization of the quarks and antiquarks created in the fragmentation process.

To answer the above-mentioned question, in a recent paper², we calculated the helicity density matrix of vector mesons by assuming that the spin of vector meson is just the sum of the spin of the quark and that of the antiquark which combine into a vector meson. By comparing the obtained results with the data, we found that the LEP data give a strong constraint on the polarization of the antiquark created in the fragmentation process. We thus extend our calculations to the deeply inelastic lepton-nucleon scattering (DIS) process and the high p_T jets in polarized high energy pp collisions. In this talk, I will make a brief summary of the calculations in Ref. 2 and 3, present the main results and make a short discussion.

*Present address: Institute of High Energy Physics, Beijing 100039, China

2. Calculation method of helicity density matrix

To calculate the helicity density matrix of a vector meson produced in the fragmentation of a polarized quark q_f^0 , we divide the produced vector mesons into the following two groups and consider them separately: (a) those which contain the initial quark q_f^0 's or the initial antiquark $\bar{q}_f^0$'s; (b) those which don't contain the initial quark or antiquark. The spin density matrix $\rho^V(z)$ for the vector meson V is given by:

$$\rho^V(z) = \sum_f \frac{\langle n(z|a, f) \rangle}{\langle n(z) \rangle} \rho^V(a, f) + \frac{\langle n(z|b) \rangle}{\langle n(z) \rangle} \rho^V(b). \quad (1)$$

Here $\langle n(z|a, f) \rangle$ and $\rho^V(a, f)$ are the average number and spin density matrix of vector mesons from (a); $\langle n(z|b) \rangle$ and $\rho^V(b)$ are those from (b). $\langle n(z) \rangle = \sum_f \langle n(z|a, f) \rangle + \langle n(z|b) \rangle$ is the total number of vector mesons and $z \equiv 2p_V/\sqrt{s}$, where p_V is the momentum of vector meson and $\sqrt{s}$ is the total e^+e^- center of mass energy. The average numbers $\langle n(z|a, f) \rangle$ and $\langle n(z|b) \rangle$ are determined by the hadronization mechanism and can be calculated using hadronization models⁴ as implemented by the Monte-Carlo events generators.

The vector mesons from group (b) are taken as unpolarized thus $\rho^V(b) = 1/3$. For those from group (a), i.e., those which contain the q_f^0 ($\bar{q}_f^0$) and a $\bar{q}$ (q) created in the fragmentation process, $V = (q_f^0 \bar{q})$ or $V = (\bar{q}_f^0 q)$, the spin density matrix $\rho^V(a, f)$ can be calculated from the direct product of the spin density matrix $\rho^{q_f^0}$ for q_f^0 (or $\rho^{\bar{q}_f^0}$ for $\bar{q}_f^0$) and that $\rho^{\bar{q}}$ for the antiquark $\bar{q}$ (or ρ^q for q). For explicity, we take $V = (q_f^0 \bar{q})$. The spin density matrix of the $q_f^0 \bar{q}$ system is $\rho^{q_f^0 \bar{q}} = \rho^{q_f^0} \otimes \rho^{\bar{q}}$. Transforming to the coupled basis $|s, s_z\rangle$ (where $\vec{s} = \vec{s}^q + \vec{s}^{\bar{q}}$), we obtain,

$$\rho^V(a, f) = \frac{2}{3 + P_f P_z} \begin{pmatrix} c_{1f}(1 + P_z) & \frac{c_{1f}}{\sqrt{2}}(P_x - iP_y) & 0 \\ \frac{c_{1f}}{\sqrt{2}}(P_x + iP_y) & \frac{1}{2}(1 - P_f P_z) & \frac{c_{2f}}{\sqrt{2}}(P_x - iP_y) \\ 0 & \frac{c_{2f}}{\sqrt{2}}(P_x + iP_y) & c_{2f}(1 - P_z) \end{pmatrix}. \quad (2)$$

where $c_{1f} = (1 + P_f)/2$, $c_{2f} = (1 - P_f)/2$ and P_f is the polarization of the initial quark q_f^0 ; $\vec{P}(P_x, P_y, P_z)$ is the polarization vector of $\bar{q}$ in the helicity frame of q_f^0 in which the z -axis is taken as the moving direction of q_f^0 .

After transforming the above result to the helicity basis of the produced vector meson, we obtain the helicity density matrix element ρ_{00} , i.e., the probability for vector meson in the helicity zero state,

$$\rho_{00}^V(a, f) \approx (1 - P_f P_z)/(3 + P_f P_z). \quad (3)$$

From this result, we can already see that $\rho_{00}^V(z, |a, f) > 1/3$ only if P_f and P_z have opposite signs. Both OPAL and DELPHI data¹ explicitly show that $\rho_{00}^V > 1/3$ for all the vector mesons except for ω . This implies that $P_z \neq 0$ and has the opposite sign as P_f in these cases.

3. Results for $e^+e^- \rightarrow VX$

We use Eq. (3) to calculate $\rho_{00}^V(a, f)$ in which the only free parameter is P_z . We first calculate ρ_{00} for K^{*0} as a function of z since the corresponding data is available¹. To fit the data, the value of P_z has to be quite large. In Fig. 2 we show the results obtained by taking $P_z=0.48$, which can fit the data reasonably well.

Similar calculations can certainly be made for other vector mesons such as ρ , ϕ , $D^{*\pm}$, B^* etc. But, for these mesons, only the average values of ρ_{00} in certain z ranges are available, and we integrate the z dependent ρ_{00} over certain z region. We found out that, for most of the vector mesons, the resulting P_z can be written as $P_z = -\alpha P_f$ with α is common for most of the q_f^0 's. The results of $\alpha=0.51$ as determined above from the data for K^{*0} can be seen in Table 1 of Ref. 2.




Fig.1. Spin density matrix element ρ_{00} of K^{*0} produced in Z^0 decay as a function of the momentum fraction z . The solid line is obtained from Eq. (3) by taking $P_z = 0.48$; the dotted line corresponds to the unpolarized case(1/3); the data are from Ref. 1.

4. Results for $lp \rightarrow l'VX$

The results we obtained for the polarization of the $\bar{q}$ in the fragmentation of the longitudinally polarized q_f^0 are direct consequences of the LEP data if we accept that the spin of the produced $V=(q_f^0\bar{q})$ is the sum of the spin of q_f^0 and that of $\bar{q}$. Since the spin effects in hadronization processes are poorly studied yet, it should be interesting to see whether they are also true in other reactions. Hence, we extend the calculations to polarized DIS processes and pp collisions.

Now, we present the calculations in $lp \rightarrow l'VX$. Here we do the calculations in the center of mass frame of the hadronic system, and replace the momentum fraction z used in $e^+e^- \rightarrow VX$ by $x_F \equiv 2p_z^*/W$, where z -axis is taken as the moving direction of the intermediate boson. Using the relation in Eq.(3) and Eq.(1), we obtain the ρ_{00}^V for DIS process,

$$\rho_{00}^V(x_F) = \sum_f \frac{1 + \alpha P_f^2}{3 - \alpha P_f^2} \frac{\langle n(x_F|a, f) \rangle}{\langle n(x_F) \rangle} + \frac{1}{3} \frac{\langle n(x_F|b) \rangle}{\langle n(x_F) \rangle}. \quad (4)$$

The P_f now is the longitudinal polarization of the outgoing struck quark. We calculate ρ_{00}^V in the current fragmentation region ($x_F > 0$) of $\mu^- p \rightarrow \mu^- VX$ at $E_\mu = 500$ GeV and the results for different combinations of beam polarization P^l and target polarization P^N are shown in Fig. 2 where $\alpha=0.5$ is used. We see that when the lepton beam is polarized, the ρ_{00}^V are much larger than 1/3 for large x_F region.

For the neutrino DIS process $\nu_\mu N \rightarrow \mu^- V X$, $P_f = -1$ (or 1) for the outgoing struck quark (antiquark), which reaches the maximum. The results of ρ_{00}^V for different mesons can be found in Ref. 3. The spin alignments for K^{*+} , $\rho^\pm$ and ρ^0 increase to about 0.6 for large x_F region.




Fig.2. Spin alignments of different vector mesons in the current fragmentation region of $\mu^- p \rightarrow \mu^- V X$ at $E_\mu = 500$ GeV. The straight line corresponds to $\rho_{00} = 1/3$.

5. Results for pp collisions

In high energy pp collisions, such as at RHIC, the vector mesons in high p_T jets can be considered as the fragmentation products of the outgoing quarks in the hard scattering subprocess. When the beam or/and target proton is longitudinally polarized, the polarization of the incoming partons can be known from the polarized distribution functions⁵ and the polarization of the outgoing partons can be calculated by perturbative QCD. Our results of spin alignments for vector mesons with $p_T > 13$ GeV at $\sqrt{s} = 500$ GeV as a function of rapidity η in beam polarized pp collisions are shown in Fig.3 where the same α is used as in previous cases.




Fig.3. Spin alignments of different vector mesons of $p_T > 13$ GeV in $\bar{p}p \rightarrow V X$ at $\sqrt{s} = 500$ GeV.

Acknowledgements

This work was supported in part by the National Science Foundation of China (NSFC) and the Education Ministry of China.

References

1. DELPHI Collab., P. Abreu *et al.*, Phys. Lett. **B406**, 271 (1997); OPAL Collab., K. Ackerstaff *et al.*, Phys. Lett. **B412**, 210 (1997); OPAL Collab., G. Abbiendi *et al.*, Eur. Phys. J. **C16**, 61 (2000), and references given therein.
2. Xu Qing-hua, Liu Chun-xiu and Liang Zuo-tang, Phys. Rev. **D63**, R111301 (2001).
3. Xu Qing-hua and Liang Zuo-tang, in preparation.
4. B. Anderson *et al.*, Phys. Rep. **97**, 31 (1983).
5. M. Glück *et al.*, Phys. Rev. **D63**, 094005 (2001).